

# Forest Vulnerability

## Project Brief

National Park Service  
U.S. Department of the Interior



Natural Resource Stewardship & Science  
Climate Change Response Program

## Climate, Trees, Pests, and Weeds: Change, Uncertainty, and Biotic Stressors at Harpers Ferry National Historical Park

### Background

Climate change and nonnative tree pests and plants affect forest composition, structure, and function, and the coming decades will likely bring accelerating changes due to these multiple global change factors. We investigated potential forest change in response to climate, differences in projections of change among climate scenarios (uncertainty), and levels of nonnative biotic stressor (tree pests and invasive plants) at 121 national park units in the eastern U.S. (see [Fisichelli et al. 2014](#) for the full article and the [project brief](#) for a synthesis). For Harpers Ferry National Historical Park, we provide here park-specific climate (Table 1) and tree habitat suitability projections (Table 2) for 2100 under two climate scenarios ("least change" and "major change"). These scenarios bracket a range of plausible future conditions based on greenhouse gas emissions and global climate model projections and facilitate assessment of uncertainty in future projections. We also report nonnative tree insects and diseases ("pests") with infestation areas that include the park (Table 3). Park-level nonnative vascular plant ("weeds") information is available from [NPSpecies](#).

Adaptation to ongoing climate change requires revising existing strategies to meet traditional goals and will increasingly require revising goals and developing new strategies. Data presented here is intended to inform managers of potential future change (climate and forests) and exacerbating factors (nonnative species).

### More Information

This project is part of ongoing work of the NPS Climate Change Response Program to support park adaptation to changing conditions (websites: for [managers](#), for the [public](#)).

### Source Publication

Fisichelli, N. A., S. R. Abella, M. P. Peters, and F. J. Krist Jr. 2014. Climate, trees, pests, and weeds: change, uncertainty, and biotic stressors in eastern U.S. national park forests. *Forest Ecology and Management* 327:31-39.

### Contact

Nicholas Fisichelli, Ph.D.  
Ecologist, Climate Change Response Program  
National Park Service  
970-225-3578  
[Nicholas.Fisichelli@nps.gov](mailto:Nicholas.Fisichelli@nps.gov)

**Table 1.** Baseline (1961-1990) and future (2070-2099) climates used in tree habitat suitability models for Harpers Ferry National Historical Park. Future projections for the two climate scenarios ("Least Change" and "Major Change") illustrate a plausible range of future conditions and are shown as departures from baseline values. The Least Change scenario includes a major decrease in future greenhouse gas emissions while the Major Change scenario is a continuation of historical increases in emissions. Climate data are averaged across a roughly 6200 mi<sup>2</sup> (16000 km<sup>2</sup>) area centered on the park.

Climate Variable	Baseline (1961-1990)	Least Change (2070-2099)	Major Change (2070-2099)
mean annual temperature	11.3 °C (52.4 °F)	+2.1 °C (+3.7 °F)	+6.9 °C (+12.5 °F)
mean January temperature	-0.7 °C (30.8 °F)	+1.7 °C (+3 °F)	+5.2 °C (+9.4 °F)
mean July temperature	23.1 °C (73.5 °F)	+2 °C (+3.6 °F)	+8.2 °C (+14.8 °F)
seasonality (July-January temp.)	23.7 °C (42.7 °F)	+0.3 °C (+0.6 °F)	+3 °C (+5.4 °F)
mean May-September temp.	20.2 °C (68.3 °F)	+2.1 °C (+3.7 °F)	+7.7 °C (+13.8 °F)
annual precipitation	1002 mm (39.5 in)	+12 %	+17 %
May-September precipitation	462 mm (18.2 in)	+10.4 %	+11.1 %

**Table 2.** Potential changes in habitat suitability (2100 compared with 1990) for tree species in Harpers Ferry National Historical Park. Species are grouped into decreasing, no change, increasing, mixed results, and new habitat groups based on change class designations for two future climate scenarios (the “least change” scenario represents strong cuts in greenhouse gas emissions and modest climatic changes and the “major change” scenario represents continued increasing greenhouse gas emissions and rapid warming). Change class designations are based on the ratio of future (2100) to baseline (1990) habitat suitability and baseline habitat values, e.g., for common species, large decrease is  $\leq 0.5$ , small decrease is  $> 0.5$  and  $\leq 0.8$ , no change is  $> 0.8$  and  $\leq 1.2$ , small increase is  $> 1.2$  and  $\leq 2.0$ , and large increase is  $> 2.0$ . Note: table below uses finer change classes ( $n=8$ ) than was used in the original article ( $n=3$ ) to designate uncertainty in habitat projections. Habitat suitability models from the US Forest Service [Climate Change Tree Atlas](#). Contact author for further details on change class definitions.

#### Decreases in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
<i>Acer pensylvanicum</i>	striped maple	extirpated	extirpated
<i>Acer rubrum</i>	red maple	small decrease	large decrease
<i>Amelanchier spp.</i>	serviceberry	small decrease	large decrease
<i>Asimina triloba</i>	pawpaw	extirpated	extirpated
<i>Betula lenta</i>	sweet birch	large decrease	large decrease
<i>Fagus grandifolia</i>	American beech	small decrease	large decrease
<i>Fraxinus americana</i>	white ash	small decrease	large decrease
<i>Liriodendron tulipifera</i>	yellow-poplar	small decrease	large decrease
<i>Magnolia acuminata</i>	cucumbertree	small decrease	small decrease
<i>Pinus rigida</i>	pitch pine	large decrease	large decrease
<i>Pinus strobus</i>	eastern white pine	large decrease	large decrease
<i>Pinus virginiana</i>	Virginia pine	small decrease	large decrease
<i>Populus grandidentata</i>	bigtooth aspen	large decrease	large decrease
<i>Prunus pensylvanica</i>	pin cherry	extirpated	extirpated
<i>Prunus serotina</i>	black cherry	large decrease	large decrease
<i>Quercus palustris</i>	pin oak	extirpated	large decrease
<i>Quercus prinus</i>	chestnut oak	small decrease	large decrease
<i>Quercus rubra</i>	northern red oak	small decrease	large decrease
<i>Robinia pseudoacacia</i>	black locust	small decrease	large decrease
<i>Tsuga canadensis</i>	eastern hemlock	large decrease	large decrease
<i>Ulmus rubra</i>	slippery elm	small decrease	large decrease

#### No Change in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
<i>Acer negundo</i>	boxelder	no change	no change
<i>Carya tomentosa</i>	mockernut hickory	no change	no change
<i>Cercis canadensis</i>	eastern redbud	no change	no change
<i>Quercus velutina</i>	black oak	no change	no change
<i>Tilia americana</i>	American basswood	no change	no change

#### Increases in Potential Habitat

Scientific Name	Common Name	Least Change	Major Change
<i>Carya ovata</i>	shagbark hickory	large increase	large increase
<i>Celtis occidentalis</i>	hackberry	small increase	large increase
<i>Diospyros virginiana</i>	common persimmon	large increase	large increase
<i>Juniperus virginiana</i>	eastern redcedar	small increase	small increase
<i>Liquidambar styraciflua</i>	sweetgum	large increase	large increase

### **Increases in Potential Habitat (continued)**

<b>Scientific Name</b>	<b>Common Name</b>	<b>Least Change</b>	<b>Major Change</b>
<i>Pinus echinata</i>	shortleaf pine	large increase	large increase
<i>Pinus taeda</i>	loblolly pine	large increase	large increase
<i>Quercus falcata var.falcata</i>	southern red oak	large increase	large increase
<i>Quercus marilandica</i>	blackjack oak	large increase	large increase
<i>Quercus muehlenbergii</i>	chinkapin oak	large increase	large increase
<i>Quercus stellata</i>	post oak	large increase	large increase

### **Mixed Results**

<b>Scientific Name</b>	<b>Common Name</b>	<b>Least Change</b>	<b>Major Change</b>
<i>Acer saccharinum</i>	silver maple	no change	large increase
<i>Acer saccharum</i>	sugar maple	no change	extirpated
<i>Carpinus caroliniana</i>	American hornbeam	small decrease	small increase
<i>Carya cordiformis</i>	bitternut hickory	no change	large increase
<i>Carya glabra</i>	pignut hickory	no change	small decrease
<i>Cornus florida</i>	flowering dogwood	no change	small decrease
<i>Fraxinus pennsylvanica</i>	green ash	no change	small increase
<i>Juglans nigra</i>	black walnut	no change	large decrease
<i>Maclura pomifera</i>	osage-orange	extirpated	large increase
<i>Morus rubra</i>	red mulberry	no change	large increase
<i>Nyssa sylvatica</i>	blackgum	no change	small decrease
<i>Ostrya virginiana</i>	eastern hophornbeam	no change	small increase
<i>Oxydendrum arboreum</i>	sourwood	small increase	small decrease
<i>Pinus pungens</i>	Table Mountain pine	large decrease	no change
<i>Platanus occidentalis</i>	sycamore	no change	small increase
<i>Populus deltoides</i>	eastern cottonwood	small decrease	large increase
<i>Quercus alba</i>	white oak	no change	small decrease
<i>Quercus coccinea</i>	scarlet oak	no change	large decrease
<i>Quercus imbricaria</i>	shingle oak	small decrease	small increase
<i>Quercus phellos</i>	willow oak	no change	large increase
<i>Salix nigra</i>	black willow	extirpated	small increase
<i>Sassafras albidum</i>	sassafras	no change	small decrease
<i>Ulmus americana</i>	American elm	no change	small increase

### **New Potential Habitat**

<b>Scientific Name</b>	<b>Common Name</b>	<b>Least Change</b>	<b>Major Change</b>
<i>Carya illinoiensis</i>	pecan	--	new entry
<i>Carya texana</i>	black hickory	new entry	new entry
<i>Celtis laevigata</i>	sugarberry	new entry	new entry
<i>Gleditsia triacanthos</i>	honeylocust	--	new entry
<i>Pinus elliottii</i>	slash pine	--	new entry
<i>Quercus falcata var.pagodifolia</i>	cherrybark oak	--	new entry
<i>Quercus macrocarpa</i>	bur oak	--	new entry
<i>Quercus nigra</i>	water oak	new entry	new entry
<i>Quercus shumardii</i>	Shumard oak	--	new entry

### New Potential Habitat (continued)

Scientific Name	Common Name	Least Change	Major Change
<i>Ulmus alata</i>	winged elm	new entry	new entry
<i>Ulmus crassifolia</i>	cedar elm	--	new entry

**Table 3.** Nonnative tree insects and diseases (“tree pests”) with infestation areas that include Harpers Ferry National Historical Park. Detection scale is the finest spatial scale at which infestation data were available for the park area. Data derived from the US Forest Service Alien Forest Pest Explorer ([AFPE](#)) Database.

Scientific Name	Common Name	Detection Scale
<i>Adelges abietis</i>	eastern spruce gall adelgid	state
<i>Adelges piceae</i>	balsam woolly adelgid	state
<i>Adelges tsugae</i>	hemlock woolly adelgid	county
<i>Agrilus planipennis</i>	emerald ash borer	county
<i>Anarsia lineatella</i>	peach twig borer	state
<i>Asterolecanium variolosum</i>	golden oak scale	state
<i>Caliroa cerasi</i>	pear sawfly	state
<i>Carulaspis juniperi</i>	juniper scale	state
<i>Caulocampus acericaulis</i>	maple petiole borer	state
<i>Ceratocystis fagacearum</i>	oak wilt	county
<i>Coleophora laricella</i>	larch casebearer	state
<i>Cronartium ribicola</i>	white pine blister rust	county
<i>Cryphonectria parasitica</i>	chestnut blight	county
<i>Cryptodiaporthe populea</i>	Dothichiza canker of poplar	state
<i>Cryptorhynchus lapathi</i>	poplar-and-willow borer	state
<i>Cyrtepistomus castaneus</i>	Asiatic oak weevil	state
<i>Diaspidiotus perniciosus</i>	San Jose scale	state
<i>Diprion similis</i>	introduced pine sawfly	county
<i>Discula destructiva</i>	dogwood anthracnose	county
<i>Dryocosmus kuriphilus</i>	chestnut gall wasp	county
<i>Epinotia nanana</i>	European spruce needleminer	state
<i>Eulecanium cerasorum</i>	calico scale	state
<i>Fenusaa pusilla</i>	birch leafminer	state
<i>Fiorinia externa</i>	elongate hemlock scale	state
<i>Homadaula anisocentra</i>	mimosa webworm	state
<i>Hylastes opacus</i>	European bark beetle	county
<i>Kaliofenusaa ulmi</i>	elm leafminer	state
<i>Lepidosaphes ulmi</i>	oystershell scale	county
<i>Lymantria dispar</i>	gypsy moth	county
<i>Neodiprion sertifer</i>	European pine sawfly	state
<i>Ophiostoma novo-ulmi</i>	Dutch elm disease	state
<i>Orchestes alni</i>	European elm flea weevil	state
<i>Otiorhynchus ovatus</i>	strawberry root weevil	county
<i>Otiorhynchus sulcatus</i>	black vine weevil	state
<i>Periphyllus lyropictus</i>	Norway maple aphid	county
<i>Phyllaphis fagi</i>	woolly beech aphid	state

**Table 3 (continued).** Nonnative tree insects and diseases (“tree pests”) with infestation areas that include Harpers Ferry National Historical Park.

<b>Scientific Name</b>	<b>Common Name</b>	<b>Detection Scale</b>
<i>Phytophthora cinnamomi</i>	littleleaf disease / phytophthora root rot	county
<i>Plagiodera versicolora</i>	imported willow leaf beetle	county
<i>Popillia japonica</i>	Japanese beetle	county
<i>Pristiphora erichsonii</i>	larch sawfly	state
<i>Pristiphora geniculata</i>	mountain-ash sawfly	state
<i>Pseudaonidia duplex</i>	camphor scale	state
<i>Rhyacionia buoliana</i>	European pine shoot moth	state
<i>Scolytus multistriatus</i>	smaller European elm bark beetle	county
<i>Scolytus schevyrewi</i>	banded elm bark beetle	county
<i>Sirococcus clavigignenti-juglandacearum</i>	butternut canker	county
<i>Taeniothrips inconsequens</i>	pear thrips	county
<i>Tomicus piniperda</i>	pine shoot beetle	county
<i>Trichiocampus viminalis</i>	poplar sawfly	state
<i>Xanthogaleruca luteola</i>	elm leafbeetle	state